

Case Report

Case report on the development of a brucellosis-related epidural abscess

Şükran Köse, Süheyla Serin Senger, Gülsün Çavdar, Sibel Yavaş

Infectious Diseases and Clinical Microbiology Clinic, Tepecik Education and Research Hospital, Izmir, Turkey

Abstract

Brucellosis is an endemic disease in developing countries. The most commonly observed complications include bone-joint involvement, particularly sacroiliitis and spondylitis. Epidural abscesses caused by brucellosis are a rare complication. We describe the case of a 33-year-old man presenting with high fever, back pain, and weakness. At physical examination, the patient was found to be paraparetic. At thoracic MRI, spondylodiscitis and epidural abscess with significant cord compression were observed. In laboratory examinations, Rose Bengal and tube agglutination tests were positive in patient's serum for brucellosis, and specific antibiotherapy was initiated. Total laminectomy was performed and the abscess was aspirated. The biopsy sample was consistent with chronic non-specific inflammation in acute abscess. No growth was detected in the abscess or blood cultures. Following surgery, medical treatment was initiated and, at six weeks' follow-up, clinical and MRI findings indicated that he had recovered. The diagnosis of spinal epidural abscess due to brucellosis should be considered among differential diagnoses in endemic regions. Early diagnosis and specific treatment are important to prevent later complications.

Key words: spinal epidural abscess; brucellosis; spondylodiscitis

J Infect Dev Ctries 2011; 5(5):403-405.

(Received 6 August 2010 - Accepted 26 November 2010)

Copyright © 2011 Kose *et al.* This is an open-access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Introduction

Brucellosis is a zoonosis caused by *Brucella* spp. which are facultative intracellular organisms transmitted by unpasteurised milk or dairy products from infected animals to humans and observed as an endemic disease among animals and humans in developing countries [1]. Common symptoms are fever, weakness, decreased appetite, back and low back pain [2]. The disease can lead to some complications, most commonly bone-joint involvement (10%-80%), particularly sacroiliitis and spondylitis [3]. When they occur, *Brucella*-related epidural abscesses generally occur in the thoracic and lumbar spine regions, but can also be found as localized abscesses in the cervical spine region; they rarely lead to spinal cord compression [4-6].

We report a case of brucellosis with thoracic spondylodiscitis and epidural abscess at the thoracic level, showing that *Brucella* spp can be the causative agent of epidural abscesses.

Case report

A 33-year-old man presented to the Infectious Diseases and Clinical Microbiology Clinic of Izmir Tepecik Education and Research Hospital in

November, 2009, with high fever, back pain and weakness, and was hospitalized for further diagnosis and treatment. From the patient's anamnesis, it was found out that the patient's complaints had started nearly two weeks before admission to our clinic. It was noted that unpasteurized cheese had been consumed about a month prior to admission. At physical examination, the patient was found to be paraparetic with grade 2/5 in the left lower limb and grade 1/5 in the right lower limb; no other pathological findings could be detected. In laboratory examinations of blood samples, the white blood cell count was 9.3 K/ μ l, sedimentation was measured as 83 mm/h, and C-reactive protein (CRP) was measured as 5.56 mg/dl. At thoracic magnetic resonance imaging (MRI), there was spondylodiscitis at T9-10 level and epidural abscess between T7-8 and T10-11 levels, while significant cord compression was observed at these levels (Figure 1). Upon determining that the Rose Bengal test was positive in the serum, together with a tube agglutination titer of 1/320, treatment was initiated with doxycycline (200 mg/day orally), rifampin (600 mg /day orally), and streptomycin (1g/day intramuscularly, for three weeks). The patient was operated on and total

Figure 1. Magnetic resonance image showing spondylodiscitis and epidural abscess with significant cord



laminectomy at T8, 9, 10, and 11 was performed; the abscess was aspirated after collecting samples for pathology and culture. Biopsy sample analysis revealed chronic non-specific inflammation in acute abscess, and no growth was detected in the culture. Following surgery the patient received six weeks of medical treatment (the aforementioned antibiotic therapy), after which he recovered (based on clinical and MRI findings).

Discussion

Epidural abscess generally represents an acute infection of the thoracolumbar region and is one of the causes of rapidly developing paraplegia [7]. It is usually diagnosed based on the development of motor deficits in the lower extremities as well as the symptoms of fever, weakness, and back pain [8].

Spinal MRI examination is the gold standard among all radiological methods of diagnosis [6]. The most common agent isolated from epidural abscess cultures is *Staphylococcus aureus*, found in 60 % of cases [9].

Brucellosis is generally associated with common symptoms such as fever, weakness, and musculoskeletal complaints such as back and low back pain [10]. Vertebral osteomyelitis, paravertebral abscesses, and especially sacroiliac infections may develop during *Brucella* infection. Spinal brucellosis may appear in the form of spondylitis, spondylodiscitis, and epidural abscess [11]. The sources of epidural abscess usually include skin, upper respiratory tract infection, and osteomyelitis of adjacent vertebrae [8]. In case of *Brucella*-related epidural abscess, the source is most probably adjacent vertebrae. Several cases of brucellosis related epidural abscess have been reported in the literature [6, 12-14]. A study by İzci *et al.* in 2005 reported that a patient who had presented with leg and sciatic pain was found to have spinal epidural abscess on MRI examination and development of *Brucella melitensis* in the blood culture [13]. There are several *Brucella*-specific assays such as microbiological culture, serological assays (*e.g.* slide or tube agglutination, Coombs test, immunocapture agglutination, Brucellacapt, immunochromatographic lateral flow, enzyme-linked immunosorbent assays and the indirect fluorescent antibody test), and molecular assays (*e.g.* polymerase chain reaction and real-time PCR), and they need to be interpreted carefully, since in some cases brucellosis can produce false negative test results [15].

Brucella abscesses generally respond to antibiotic treatment [16]. Surgical treatment should be considered if there are neurological deficits resulting from nerve root or spinal cord compression [17]. In our case, there was significant cord compression and paraparesis which developed secondary to brucellosis in the lower extremities; therefore, immediate surgery was arranged for the patient. Since short-term monotherapy has a high rate of relapse (30%), generally combination therapies are preferred for the treatment of brucellosis [18-19]. The most commonly administered therapy is combination treatment with doxycycline, rifampin and streptomycin. In cases of perineural abscess, the treatment can be extended up to 6 to 12 months [20].

In conclusion, in cases of non-specific symptoms, including back and lower back pain and progressive paraparesis, the diagnosis of spinal epidural abscess

should be considered among differential diagnoses, and *Brucella* bacteria should be considered among the causative agents in endemic regions. If there are surgical indications in epidural abscesses, the abscesses should be drained as soon as possible. Otherwise, neurological deficits may develop which may become persistent.

References

- Mantur BG and Amarnath SK (2008) Brucellosis in India - a review. *J Biosci* 33: 539-547.
- Akhvlediani T, Clark DV, Chubabria G, Zenaishvili O, Hepburn MJ (2010) The changing pattern of human brucellosis: clinical manifestations, epidemiology, and treatment outcomes over three decades in Georgia. *BMC Infect Dis* 10: 346.
- Colmenero JD, Reguera JM, Martos F Sánchez-De-Mora D, Delgado M, Causse M, Martín-Farfán A, Juárez C (1996) Complications associated with *Brucella melitensis* infection: a study of 530 cases. *Medicine (Baltimore)* 75: 195-211.
- Mousa AR, Muhtaseb SA, Al mudallal DS, Khodeir SM, Marafie AA. Osteoarticular complications of brucellosis: a study (1987) of 169 cases. *Rev Infect Dis* 9: 531-543.
- Ceviker N, Baykaner K, Goksel M, Sener L, Alp H (1989) Spinal cord compression due to *Brucella* granuloma. *Infection* 17: 304-305.
- Gorgulu A, Albayrak BS, Gorgulu E, Görgülü A, Albayrak BS, Görgülü E, Tural O, Karaaslan T, Oyar O, Yilmaz M (2006) Spinal epidural abscess due to *Brucella*. *Surg Neurol* 66: 141-146; discussion 6-7.
- Gul HC, Erdem H, Bek S (2009) Overview of neurobrucellosis: a pooled analysis of 187 cases. *Int J Infect Dis* 13: e339-e343.
- Reihsaus E, Waldbaur H, Seeling W (2000) Spinal epidural abscess: a meta-analysis of 915 patients. *Neurosurg Rev* 23: 175-204; discussion 5.
- Sendi P, Bregenzer T, Zimmerli W (2008) Spinal epidural abscess in clinical practice. *QJM* 101: 1-12.
- Cokca F, Meco O, Arasil E, Unlu A (1994) An intramedullary dermoid cyst abscess due to *Brucella abortus* biotype 3 at T11-L2 spinal levels. *Infection* 22: 359-360.
- Mousa AM, Bahar RH, Araj GF, Koshy TS, Muhtaseb SA, al-Mudallal DS, Marafie AA (1990) Neurological complications of brucella spondylitis. *Acta Neurol Scand* 81: 16-23.
- Daglioglu E, Bayazit N, Okay O, Dalgic A, Hatipoglu HG, Ergunor F (2009) Lumbar epidural abscess caused by brucella species: report of two cases. *Neurocirugia (Astur)* 20: 159-162.
- Izci Y (2005) Lumbosacral spinal epidural abscess caused by *Brucella melitensis*. *Acta Neurochir (Wien)* 147: 1207-1209; discussion 9.
- Paz JF, Alvarez FJ, Roda JM, Frutos R, Isla A (1994) Spinal epidural abscess caused by *Brucella*: case report. *J Neurosurg Sci* 38: 245-249.
- Araj GF (2010) Update on laboratory diagnosis of human brucellosis. *Int J Antimicrob Agents* 36 Suppl 1: S12-17.
- Pina MA, Modrego PJ, Uroz JJ, Cobeta JC, Lerin FJ, Baiges JJ (2001) Brucellar spinal epidural abscess of cervical location: report of four cases. *Eur Neurol* 45: 249-253.
- Coskun E, Suzer T, Yalcin N, Tahta K (1998) Spinal extradural compression caused by granuloma of brucellosis. *Scand J Infect Dis* 30: 311-313.
- Bingol A, Yucemen N, Meco O (1999) Medically treated intraspinal "Brucella" granuloma. *Surg Neurol* 52: 570-576.
- Ariza J, Bosilkovski M, Cascio A, Colmenero JD, Corbel MJ, Falagas ME, Memish ZA, Roushan MR, Rubinstein E, Sipsas NV, Solera J, Young EJ, Pappas G; International Society of Chemotherapy; Institute of Continuing Medical Education of Ioannina (2007) Perspectives for the treatment of brucellosis in the 21st century: the Ioannina recommendations. *PLoS Med* 4: e317.
- Bodur H, Erbay A, Colpan A, Akinci E (2004) Brucellar spondylitis. *Rheumatol Int* 24: 221-226.

Corresponding author

Gülsün Çavdar
 İzmir Tepecik Eğitim ve Araştırma Hastanesi
 Enfeksiyon Hastalıkları ve Klinik Mikrobiyoloji Kliniği, 35110
 Yenisehir, İZMİR, TURKEY
 Telephone: +90 232 469 6969-1719
 Fax: +90 232 433 0756
 Email: gakincioglu@hotmail.com

Conflict of interests: No conflict of interests is declared.