

Coronavirus Pandemic

COVID-19, frailty and long-term care: Implications for policy and practice

Melissa K Andrew^{1,2}, Samuel D Searle^{1,3}, Janet E McElhaney⁴, Shelly A McNeil^{5,2}, Barry Clarke⁶, Kenneth Rockwood¹, David J Kelvin^{7,8,2}

¹ Department of Medicine (Geriatrics), Dalhousie University, Halifax, Canada

² Canadian Centre for Vaccinology, Dalhousie University, Halifax, Canada

³ MRC Unit for Lifelong Health and Ageing, University College London, United Kingdom

⁴ Health Sciences North Research Institute, Sudbury, Canada

⁵ Department of Medicine (Infectious Diseases), Dalhousie University, Halifax, Canada

⁶ Department of Family Medicine, Dalhousie University, Halifax, Canada

⁷ Department of Microbiology and Immunology, Dalhousie, Halifax, Canada

⁸ Laboratory of Immunity, Shantou University Medical College, Shantou, China

Abstract

Older adults have been disproportionately affected by the COVID-19 pandemic, with many outbreaks occurring in Long Term Care Facilities (LTCFs). We discuss this vulnerability among LTCF residents using an ecological framework, on levels spanning from the *individual to families and caregivers, institutions, health services and systems, communities, and contextual* government policies. Challenges abound for fully understanding the burden of COVID-19 in LTCF, including differences in nomenclature, data collection systems, cultural differences, varied social welfare models, and (often) under-resourcing of the LTC sector. Registration of cases and deaths may be limited by testing capacity and policy, record-keeping and reporting procedures. Hospitalization and death rates may be inaccurate depending on atypical presentations and whether or not residents' goals of care include escalation of care and transfer to hospital. Given the important contribution of frailty, use of the Clinical Frailty Scale (CFS) is discussed as a readily implementable measure, as are lessons learned from the study of frailty in relation to influenza. Biomarkers hold emerging promise in helping to predict disease severity and address the puzzle of why some frail LTCF residents are resilient to COVID-19, either remaining test-negative despite exposure or having asymptomatic infection, while others experience the full range of illness severity including critical illness and death. Strong and coordinated surveillance and research focused on LTCFs and their frail residents is required. These efforts should include widespread assessment of frailty using feasible and readily implementable tools such as the CFS, and rigorous reporting of morbidity and mortality in LTCFs.

Key words: COVID-19; frailty; older adult; long-term care; biomarker.

J Infect Dev Ctries 2020; 14(5):428-432. doi:10.3855/jidc.13003

(Received 09 May 2020 – Accepted 16 May 2020)

Copyright © 2020 Andrew *et al.* This is an open-access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Around the world, older adults are being hit hard by COVID-19. Outbreaks in Long-Term Care Facilities (LTCFs) are particularly devastating [1,2]. Given shortages of both people and resources, there is a pressing need to provide well-resourced and compassionate care for COVID-19 illness during outbreaks in LTCFs. Residents of LTCFs are at great risk for poor outcomes for many reasons, discussed here within an ecological framework. These multi-level factors have a major impact on the spread and severity of COVID-19, and are critical to understand if we are to achieve better outcomes in this vulnerable sector. In this paper we aim to discuss both challenges to gaining full understanding of the impact of COVID-19 in LTCF settings and the importance of frailty, an easily ascertained clinical factor, to inform risk and care

planning. We draw from related experience in the study of frailty in relation to influenza, and suggest future directions for COVID-19 research through a frailty lens.

Why have LTCF emerged as the centre of the global pandemic?

The many factors that put residents of LTCFs at great risk for poor outcomes can be conceptualized at several levels within an ecological framework (Figure 1) [3,4]. At an *individual* level, LTCF residents have many chronic conditions, contributing to their high levels of frailty [5]. One consequence is that any disease, including COVID-19, presents differently (“atypically”) in them [6]. This is a pragmatic challenge to case definition and ascertainment, making them more

vulnerable to severe outcomes. Cases going unrecognized have clinical and infection control ramifications, and increases the risk of future transmission. At the level of *family, friends and peer groups*, visitor limitations aim to limit introduction of illness into the facility but also serve to isolate residents from those who could be advocates (e.g. in the detection of early signs that “something is not right” as subtle signs of illness begin). On an *institutional* facility level, prior to the COVID-19 pandemic, social interaction and sense of community within facilities were considered something to strive for. These are now considered as contributors of spread of COVID-19 within facilities, along with others facility characteristics including close living quarters and shared dining facilities. Limited resources (and sometimes motives for profit) contribute to staffing models that rely on low paid staff without benefits and thus unable to take sick leave, or inadvertently spreading infection among facilities by working part-time at several LTCF. At the level of *communities*, disease transmission may be facilitated when facilities and/or staff are located in community clusters or marginalized communities. Relationships with community planning also arise, for example when staff must rely on crowded public transit (or transit routes and schedules that are affected by pandemic measures taken by transit authorities) to commute to and between their workplaces. On a *contextual policy level* within jurisdictions, funding and resourcing may be suboptimal for care provision at the best of times, even without the added demands of local outbreaks within a global pandemic. There may be barriers to testing due to limited resources or other constraints or agendas.


These factors have a major impact on the spread and severity of COVID-19, and are critical to understand if we are to achieve better outcomes in this vulnerable sector.

Challenges to understanding the impact of COVID-19 in LTCFs

Reliable estimates of the impact of COVID-19 on morbidity and mortality within LTCFs are hard to come by. In Canada, it has been reported that 79% of deaths have been in LTCFs [7]. In Europe, deaths in care homes reportedly account for approximately a third to more than a half of COVID-19 deaths across several countries [8].

There are many challenges to achieving a full understanding of the full implications of COVID-19 in this population. LTCFs come in all shapes and sizes, are called different names in different jurisdictions (e.g. care homes, retirement communities, Skilled Nursing Facilities, convalescent homes, homes for the aged etc...), and may or may not be formally regulated or registered with local authorities and data collection systems. Cultural differences (e.g. the common Asian doctrine of filial piety) [9], varied social welfare models (e.g. expectations on family vs. state for care of older adults) [10] and differential resource availability also impact the numbers and organization of such residences. When it comes to registering cases of COVID-19 and the morbidity and mortality the SARS-CoV-2 virus leaves in its wake, additional layers of challenges abound. Testing capacity and policy are understandably highly variable. Hospitalization rates may be misleading as an indicator of illness burden case mix severity where care pathways exist to support managing illness in place within the LTCF. Importantly, LTCF residents with severe illness whose goals of care do not include transfer to acute care or further escalation of care (e.g. mechanical ventilation, Intensive Care Unit) may stay in LTCF and thus not get counted [11]. Residents who presented atypically and did not get tested before their death will also go uncounted, or COVID-19 may not be reported as a cause of death if a frail resident was seen to die “with” rather than “of” COVID-19. Analyses based on place of

Figure 1. Ecological model of COVID-19’s impact on LTCFs.



death recorded on death certificates will exclude any residents who have been transferred to hospital and die there.

Frailty, vulnerability and resilience

Frailty, which is a measure of how the accumulation of health and functional problems makes older people vulnerable to poor health outcomes, is an important part of this story [12]. It is imperative to better understand how clinical assessments of frailty, ideally integrated with routinely collected clinical data, can be used as predictors of illness severity and recovery. Many measures of frailty exist, from those based on deficit accumulation abstracted from Electronic Medical Record data or from comprehensive geriatric assessments, to performance-based measures or a frailty phenotype, to integrative measures based on clinical judgment [13]. The right tool for the job will of course vary by setting and according to existing data sources. When considering how to efficiently capture frailty status in pandemic management settings where time and resources are limited, feasibility and ease of use are paramount. The Clinical Frailty Scale has emerged as a widely used measure for these reasons. The CFS is a 9 category scale based on a clinician's judgment (or even self- or caregiver report) in which a patient is categorized on a scale where 1 = very fit, 2 = well, 3 = managing well with treated comorbid illness, 4 = vulnerable, 5 = mild frailty, 6 = moderate frailty, 7 = severe frailty, 8 = very severe frailty, and 9 = terminally ill. (<https://www.dal.ca/sites/gmr/our-tools/clinical-frailty-scale.html>) It has been well validated and used in more than 20 countries, and numerous translations are available including a freely downloadable App for use on mobile devices [14,15].

Notably, the story is not simply one of understanding and grading *vulnerability* - it is also critical to understand *resilience*, as in why it is that some frail LTCF residents end up with mild (or no) symptoms, while others, just as frail, become critically ill [16]. Needed is an understanding of what factors lead to higher risk for more severe illness, and how to maximize residents' chances of recovering with good quality of life.

Lessons learned from frailty and influenza

Because of its importance for influencing risk and outcomes of infections such as influenza and COVID-19, frailty is ideally integrated into surveillance efforts. For example, the Canadian Immunization Research Network's Serious Outcomes Surveillance (SOS) Network has long included a focus on measures of

health status and outcomes that are relevant for older adults, notably frailty. Key findings have included: a) the importance of frailty for understanding vaccine effectiveness [17], b) that frailty is associated with atypical illness presentation and the potential for under-recognition of influenza cases using ILI case definitions [18], c) baseline frailty sets older adults up for poor recovery following infection (including further progression of frailty itself) [19], and d) hospitalization with influenza not uncommonly leads to an important burden of persistent functional decline and need for increased caregiving supports including incident LTCF placement [20]. As international recognition of the importance of frailty mounts, the Clinical Frailty Scale has been included in data collection of the Global Influenza Hospital Surveillance Network (<https://www.gihsn.org/the-network/protocol>), for use in sites around the world in low, middle and high income country settings.

Can biomarkers predict COVID-19 severity?

A search is under way to identify biomarkers and immune function indicators to try and understand how older adults' immune responses help (or hinder) in fighting off the illness. These include biomarkers associated with inflammation, endothelial function, mitochondria and apoptotic function, calcium homeostasis, fibrosis, neuromuscular function, sarcopenia, and bone/hormone metabolism and nutritional status [21,22]. Again drawing from experience on influenza, while it is well known that older people are not well-protected by current influenza vaccines, what is less understood is that frailty is an important predictor of vaccine effectiveness against influenza-related hospitalization [17]. However, there are limited studies of the relationship between frailty and immune biomarkers of the response to influenza vaccination, [23-25] influenza illness, or other respiratory infections such as COVID-19 [26]. Monitoring cytokine levels in the plasma or serum has yielded important clues to disease severity in SARS-CoV-1, pandemic influenza (H1N1) and COVID-19 infections [27-29]. Elevated levels of IL-6 is a biomarker for severe disease in, influenza and COVID-19 and IL-6 levels are used to guide treatment with IL-6 receptor blockers [27,28]. The interest in biomarkers is that 1) along with clinical history biomarkers may be useful in identifying individuals at the highest risk for severe COVID-19 infection, 2) biomarkers may be used to stratify patients into treatment groups, and 3) biomarkers may yield important information on the mechanisms of severe disease.

Just as it is not clear how frailty affects illness presentation and recovery, it is not clear whether similar immune responses will be shared in healthy individuals versus older frail residents of LTCF. It is therefore imperative that older frail participants are included in studies so that findings from healthier people are not generalized to the older, frailer and more high-risk population.

Implications for policy and practice

An understanding of frailty and the contextual factors outlined in the ecological model presented above is important to inform clinical and policy decision-making on how care and management strategies can be implemented to best support COVID-19 outbreaks. Understanding frailty also obliges an acknowledgement of the many medical and social factors that contribute to illness in older people. It underscores why so much care provision, modeled on people having a single dominant illness at a time, has such difficulty in managing older people with “the ‘flu’”, resulting in increased risk of admission to long term care [20].

Strong and coordinated surveillance and research focused on LTCFs and their frail residents is required to provide fundamental insights as to how COVID-19 affects this high risk and vulnerable population, so that we may develop and inform efforts to develop better prevention and treatment strategies to combat COVID-19. Addressing locally relevant factors at each of the levels of influence presented in the ecological framework will be important. We recommend that these efforts should include widespread assessment of frailty using feasible and easily implementable tools such as the CFS. Rigorous reporting of morbidity and mortality in LTCFs is also required so that residents, families and the general public can be reassured that all efforts are being made to provide well-resourced and compassionate care for residents of LTCFs during the COVID-19 pandemic and beyond.

Acknowledgements

This work was supported by the Li-Ka Shing Foundation, Shantou University Medical College; a Rapid Response award for COVID-19, Canadian Institutes of Health Research; Dalhousie Medical Research Foundation; and a Rapid Response Award from Research Nova Scotia. DJK holds the Canada Research Chair in Translational Vaccinology and Inflammation at Dalhousie University.

References

1. McMichael TM, Currie DW, Clark S, Pogojans S, Kay M, Schwartz NG, Lewis J, Baer A, Kawakami V, Lukoff MD, Ferro J, Brostrom-Smith C, Rea TD, Sayre MR, Riedo FX, Russell D, Hiatt B, Montgomery P, Rao AK, Chow EJ, Tobolowsky F, Hughes MJ, Bardossy AC, Oakley LP, Jacobs JR, Stone ND, Reddy SC, Jernigan JA, Honein MA, Clark TA, Duchin JS (2020) Epidemiology of Covid-19 in a long-term care facility in King County, Washington. *N Engl J Med* 382: 2005-2011.
2. D'Adamo H, Yoshikawa T, Ouslander JG (2020) Coronavirus disease 2019 in geriatrics and long-term care: The ABCDs of COVID-19. *J Am Geriatr Soc* 68: 912-917.
3. Andrew MK, Keefe J (2014) Social vulnerability among older adults: a social ecology perspective from the National Population Health Survey of Canada. *BMC Geriatrics* 14: 90.
4. Andrew MK, Dupuis-Blanchard S, Maxwell C, Giguere A, Keefe J, Rockwood K, St John P (2018) Social and societal implications of frailty, including impact on Canadian healthcare systems. *J Frailty Aging* 7: 217-223.
5. Theou O, Tan EC, Bell JS, Emery T, Robson L, Morley JE, Rockwood K, Visvanathan R. (2016) Frailty levels in residential aged care facilities measured using the Frailty Index and FRAIL-NH scale. *J Am Geriatr Soc* 64: e207-e212.
6. Cesari M, Montero-Odasso M (2020) COVID-19 and older adults. Lessons learned from the Italian epicenter. *Can Geriatr J*: 23: 152-154.
7. Walsh M, Semeniuk I (2020) Long-term care connected to 79 per cent of COVID-19 deaths in Canada. <https://www.theglobeandmail.com/politics/article-long-term-care-connected-to-79-per-cent-of-covid-19-deaths-in-canada/>. The Globe and Mail, 28 April, 2020, Ottawa and Toronto.
8. Comas-Herrera A, Zalakaín J, Litwin C, Hsu AT, Lane N, Fernández JL (2020) Mortality associated with COVID-19 outbreaks in care homes: early international evidence. Available: <https://ltccovid.org/2020/05/04/updated-report-mortality-associated-with-covid-19-outbreaks-in-care-homes-early-international-evidence/>. Accessed: 25 May 2020.
9. Chen L (2017) Power and ambivalence in intergenerational communication: Deciding to institutionalize in Shanghai. *J Aging Stud* 41: 44-51.
10. Wallace L, Theou O, Pena F, Rockwood K, Andrew MK (2014) Social vulnerability as a predictor of mortality and disability: Cross-country differences in the Survey of Health, Aging, and Retirement in Europe (SHARE). *Aging Clin Exp Res* 27: 365-372
11. Nemiroff L, Marshall EG, Jensen JL, Clarke B, Andrew MK (2019) Adherence to "No transfer to hospital" advance directives among nursing home residents. *J Am Med Dir Assoc* 20: 1373-1381.
12. Clegg A, Young J, Iliffe S, Rikkert MO, Rockwood K (2013) Frailty in elderly people. *Lancet* 381: 752-762.
13. Theou O, Brothers TD, Mitnitski A, Rockwood K (2013) Operationalization of frailty using eight commonly used scales and comparison of their ability to predict all-cause mortality. *J Am Geriatr Soc* 61: 1537-1551.
14. Rockwood K, Song X, MacKnight C, Bergman H, Hogan DB, McDowell I, Mitnitski A. (2005) A global clinical measure of fitness and frailty in elderly people. *CMAJ* 173: 489-495.
15. Dalhousie Geriatric Medicine Research (2008) Clinical Frailty Scale Available: <https://www.dal.ca/sites/gmr/our-tools/clinical-frailty-scale.html>. Accessed: 8 May 2020.

16. Kimball A, Hatfield KM, Arons M, James A, Taylor J, Spicer K, Bardossy AC, Oakley LP, Tanwar S, Chisty Z, Bell JM, Methner M, Harney J, Jacobs JR, Carlson CM, McLaughlin HP, Stone N, Clark S, Brostrom-Smith C, Page LC, Kay M, Lewis J, Russell D, Hiatt B, Gant J, Duchin JS, Clark TA, Honein MA, Reddy SC, Jernigan JA, Public Health S, King C, Team CCI (2020) Asymptomatic and presymptomatic SARS-CoV-2 infections in residents of a long-term care skilled nursing facility - King County, Washington, March 2020. *Morb Mortal Wkly Rep* 69: 377-381.
17. Andrew MK, Shinde V, Ye L, Hatchette T, Haguinet F, Dos Santos G, McElhaney JE, Ambrose A, Boivin G, Bowie W, Chit A, ElSherif M, Green K, Halperin S, Ibaruchi B, Johnstone J, Katz K, Langley J, Leblanc J, Loeb M, MacKinnon-Cameron D, McCarthy A, McGeer A, Powis J, Richardson D, Semret M, Stiver G, Trottier S, Valiquette L, Webster D, McNeil SA, Serious Outcomes Surveillance Network of the Public Health Agency of Canada/Canadian Institutes of Health Research Influenza Research N, the Toronto Invasive Bacterial Diseases N (2017) The importance of frailty in the assessment of influenza vaccine effectiveness against influenza-related hospitalization in elderly people. *J Infect Dis* 216: 405-414.
18. Andrew MK, McElhaney JE, McGeer AA, Hatchette TF, Leblanc J, Webster D, Bowie W, Poirier A, Nichols MK, McNeil SA, Canadian Immunization Research Network Serious Outcomes Surveillance Network I (2020) Influenza surveillance case definitions miss a substantial proportion of older adults hospitalized with laboratory-confirmed influenza: A report from the Canadian Immunization Research Network (CIRN) Serious Outcomes Surveillance (SOS) Network. *Infect Control Hosp Epidemiol* 41: 499-504.
19. Lees C, Godin J, McElhaney JE, McNeil SA, Loeb M, Hatchette TF, LeBlanc J, Bowie W, Boivin G, McGeer A, Poirier A, Powis J, Semret M, Webster D, Andrew MK (2020) Frailty hinders recovery from influenza and acute respiratory illness in older adults. *J Infect Dis* jiaa092: Online ahead of print
20. Godin J, Theou O, Black K, McNeil SA, Andrew MK (2019) Long-term care admissions following hospitalization: The role of social vulnerability. *Healthcare* 7: 91.
21. Guest PC. (2020) The Impact of New Biomarkers and Drug Targets on Age-Related Disorders. *Methods Mol Biol.* 2138: 3-28.
22. Marron MM, Harris TB, Boudreau RM, Clish CB, Moore SC, Murphy RA, Murthy VL, Sanders JL, Shah RV, Tseng GC, Wendell SG, Zmuda JM, Newman AB (2020) A metabolite composite score attenuated a substantial portion of the higher mortality risk associated with frailty among community-dwelling older adults. *J Gerontol A Biol Sci Med Sci* glaa112: Online ahead of print.
23. Loeb N, Andrew MK, Loeb M, Kuchel GA, Haynes L, McElhaney JE, Verschoor CP. Frailty is associated with increased hemagglutinin-inhibition titres in a 4-year randomized trial comparing standard and high dose influenza vaccination. *Open Forum Infect Dis.* In press.
24. DiazGranados CA, Dunning AJ, Robertson CA, Talbot HK, Landolfi V, Greenberg DP. (2015) Efficacy and immunogenicity of high-dose influenza vaccine in older adults by age, comorbidities, and frailty. *Vaccine* 33: 4565-4571.
25. Keshtkar-Jahromi M, Ouyang M, Keshtkarjahromi M, Almed S, Li H, Walston JD, Rios R, Leng SX (2018) Effect of influenza vaccine on tumor necrosis factor-like weak inducer of apoptosis (TWEAK) in older adults. *Vaccine* 36: 2220-2225.
26. Johnstone J, Parsons R, Botelho F, Millar J, McNeil S, Fulop T, McElhaney J, Andrew MK, Walter SD, Devereaux PJ, Malekesmaeili M, Brinkman RR, Mahony J, Bramson J, Loeb M (2014) Immune biomarkers predictive of respiratory viral infection in elderly nursing home residents. *PLoS One* 9: e108481.
27. Paquette SG, Banner D, Zhao Z, Fang Y, Huang SS, Leomicronn AJ, Ng DC, Almansa R, Martin-Loeches I, Ramirez P, Socias L, Loza A, Blanco J, Sansonetti P, Rello J, Andaluz D, Shum B, Rubino S, de Lejarazu RO, Tran D, Delogu G, Fadda G, Kraiden S, Rubin BB, Bermejo-Martin JF, Kelvin AA, Kelvin DJ (2012) Interleukin-6 is a potential biomarker for severe pandemic H1N1 influenza A infection. *PLoS One* 7: e38214.
28. Jamilloux Y, Henry T, Belot A, Viel S, Faucher M, El Jammal T, Walzer T, Francois B, Seve P. (2020) Should we stimulate or suppress immune responses in COVID-19? Cytokine and anti-cytokine interventions. *Autoimmun Rev* 102567: Online ahead of print.
29. Cameron MJ, Bermejo-Martin JF, Danesh A, Muller MP, Kelvin DJ. (2008) Human immunopathogenesis of severe acute respiratory syndrome (SARS). *Virus Res* 133: 13-19.

Corresponding author

Melissa K Andrew
 c/o Division of Geriatric Medicine
 Dalhousie University
 5955 Veterans' Memorial Lane
 Halifax NS, Canada
 Tel +1 902 473-4995
 Fax +1 902 473-7133
 Email: mandrew@dal.ca

Conflict of interests: MKA and SAM report grant funding from the Public Health Agency of Canada through Service Canada, Canadian Institutes of Health Research, and Canadian Immunization Research Network in the conduct of the Serious Outcomes Surveillance Network. MKA reports funding and honoraria from the Canadian Frailty Network, Sanofi, GSK, and Pfizer outside the submitted work. JEM's institution has received honoraria for consultancies with GSK, Sanofi, Merck, Pfizer, Medicago and RestorBio, outside of the submitted work. SAM reports funding and honoraria from GSK, Merck, Pfizer and Sanofi outside the submitted work. KR founded and is Chief Science Officer of DGI Clinical, which has several contracts with pharma and medical device manufacturers. He developed the CFS and has asserted copyright through Dalhousie University; it is made available for free clinical and academic use including through a downloadable App. SDS, BC and DJK report no conflict of interest..